

Simple dynamics on the brane.

Marek Szydlowski

Astronomical Observatory, Jagiellonian University
Orla 171, 30-244 Kraków, Poland*

Mariusz P. Dąbrowski

Institute of Physics, University of Szczecin
Wielkopolska 15, 70-451 Szczecin, Poland†

Adam Krawiec

Institute of Public Affairs, Jagiellonian University
Rynek Główny 8, 31-042 Kraków, Poland‡

February 1, 2008

Abstract

We apply methods of dynamical systems to study the behaviour of the Randall-Sundrum models. We determine evolutionary paths for all possible initial conditions in a 2-dimensional phase space and we investigate the set of accelerated models. The simplicity of our formulation in comparison to some earlier studies is expressed in the following: our dynamical system is a 2-dimensional Hamiltonian system, and what is more advantageous, it is free from the degeneracy of critical points so that the system is structurally stable. The phase plane analysis of Randall-Sundrum models with isotropic Friedmann geometry clearly shows that qualitatively we deal with the same types of evolution as in general relativity, although quantitatively there are important differences.

PACS number(s): 04.20.Jb, 04.65.+e, 98.80.-k, 98.80.Hw

*email: uoszydlo@cyf-kr.edu.pl

†email: mpdabfz@uoo.univ.szczecin.pl

‡email: uukrawie@cyf-kr.edu.pl

1 Introduction

The rapid development of particle-physics-motivated cosmology represented mainly by superstring cosmology [1, 2, 3] has led to a change of views onto the standard problems of inflationary cosmology such as the (past) horizon and flatness problems [4, 5]. On the other hand, the astronomical observations of supernovae Ia [6] strongly suggest that the universe not only had possibly accelerated at its early stages of evolution but it is also accelerating now. This puts strong constraints onto the matter content of the universe because only the exotic (negative pressure) matter in standard cosmology can lead to acceleration. However, a phenomenological nature of such an exotic matter in standard models expressed in terms of the perfect fluid does not seem to make enough connection with particle physics and that is why one is looking for other, more physical, descriptions. Among them the main proposal is quintessence or time-dependent scalar field [7] which substitutes an ordinary phenomenological barotropic fluid. However, there are other interesting proposals to express the phenomenon of acceleration which are related to superstring or M-theory models. The best known are pre-big-bang [1, 2], brane [8, 9, 10] and ekpyrotic models [11, 12]. In this paper we try to study the standard cosmology problems (such as cosmic acceleration and past horizon problems) within the framework of the brane universes. We do not study, for instance, the future horizon problem following the recent discussion inspired by the S-matrix formulation within the superstring theories [14]. In the paper we are interested mainly in the early stage of the evolution of the universe although the supernovae data gives restrictions onto this stage and they should be taken into account.

The idea of brane universes has been first presented by Hořava and Witten [8] who considered strong coupling limit of heterotic $E_8 \times E_8$ superstring theory, i.e., M-theory. This limit results in ‘exotic’ [15, 16] Kaluza-Klein type compactification of $N = 1$, $D = 11$ supergravity on a S^1/Z_2 orbifold (a unit interval) in a similar way as compactification of $N = 1$, $D = 11$ supergravity on a circle S^1 results in strongly coupled limit of type IIA superstring theory. In Hořava-Witten theory there exist two 10-dimensional branes to which all the gauge interactions are confined, and they are connected via the orbifold, with gravity propagating in all 11 dimensions. After further compactification of Hořava-Witten models on a Calabi-Yau manifold one gets an effective 5-dimensional theory which has been applied to cosmology [17, 18, 19, 20].

Randall and Sundrum [9, 10] developed similar to Hořava-Witten scenario which was mainly motivated by the hierarchy problem in particle physics [21, 23]. As a result, they obtained a 5-dimensional spacetime (bulk) with Z_2 symmetry with two/one 3-brane(s) embedded in it to which all the gauge

interactions are confined. In one-brane scenario [10] the brane appears at the $y = 0$ position, where y is an extra dimension coordinate and the 5-dimensional spacetime is an anti-de Sitter space with negative 5-dimensional cosmological constant. The extra dimension can be infinite due to the exponential ‘warp’ factor in the metric

$$ds^2 = \exp\left(-2\frac{|y|}{l}\right) [-dt^2 + d\vec{x}^2] + dy^2, \quad (1)$$

and l gives the curvature scale of the anti-de Sitter space. In the simplest case the induced metric on a brane is a Minkowski metric (energy momentum tensor of matter vanishes). However, the requirement to allow matter energy-momentum tensor on the brane leads to breaking of conformal flatness in the bulk, and the metric (1) is no longer valid. This fact is obviously related to the appearance of the Weyl curvature in the bulk [24, 25, 30]. The full set of 5-dimensional and projected 4-dimensional equations has been presented in Refs. [26, 27, 28]. Global geometric properties of such brane models have also been studied (see Ref. [29]). Generalized bulk spacetimes different from those of (1) have been found in Ref. [31]. Anisotropic Kasner branes have been immersed in AdS bulk in Ref. [32].

Campos and Sopena [33, 34] used the dynamical system methods to the analysis of the Friedmann, Bianchi I and V Randall-Sundrum brane world type cosmological models with a non-vanishing 5-dimensional Weyl tensor. They considered the dynamics of this model in the form of a *higher-than-two-dimensional* dynamical system. Exact analytic brane configurations with a vanishing Weyl tensor with perfect and viscous fluid have been presented in Refs. [35, 36]. Coley [37, 38] also studied the dynamics of these Randall-Sundrum models - he made a step towards Mixmaster (Bianchi IX) dynamics and found it was not chaotic provided the matter had positive pressure. In this paper we show that the dynamical system which describes the evolution of the brane models (both isotropic (FRW) and anisotropic (Bianchi I or Bianchi V) types) can be represented in the simplest way in the form of a *two-dimensional* Hamiltonian dynamical system. Such visualization has a great advantage because we *avoid* the problem of degeneracy of critical points which appeared in the *higher dimensional* phase space (see Ref. [13] in [34]). It is well known that the existence of such critical points is a possible reason for the structural instability of a model. On the other hand, the representation of dynamics as a one-dimensional Hamiltonian flow allows to make the classification of possible evolution paths in the *configuration space* which is complementary to phase diagrams. It also makes simpler to discuss the physical content of the model. Finally, the construction of the Hamiltonian allows to study quantum cosmology on the brane as it was

attempted in Refs. [39, 40], in full analogy to what is usually done in general relativity [41].

In this paper we demonstrate the effectiveness of representation of dynamics as a one-dimensional Hamiltonian flow. In this representation the phase diagrams in a two-dimensional phase space allow to analyze the *acceleration and (past) horizon* problems in a clear way. We reduce the dynamics to a two-dimensional phase space with an autonomous system of equations.

We deal with the full global dynamics of brane models whose asymptotic states are represented by critical points of the system. From theoretical point of view it is important how large the class of accelerated models is. We will call this class of accelerated models typical/generic, if the domain of acceleration in the phase space is a non-zero measure. On the other hand, if only the non-generic (zero-measure) trajectories are represented by accelerated models, then the mechanism which drives these trajectories should be called ineffective. Such a point of view is a consequence of the fact that, if the acceleration is an attribute of a trajectory which starts with a given initial conditions, it should also be an attribute of a trajectory which starts with a neighbouring initial conditions.

Our analysis of the brane-world type model may be considered as complementary to Campos and Sopuerta's analysis [33, 34] (see also [37, 38]). However, it is more advantageous in many points. It is because our dynamical system is a 2-dimensional system which is the smallest possible dimension to study isotropic cosmological systems of equations. This allows to avoid huge redundancy of degenerated critical points and trajectories which appear in a higher dimensional phase space (cf. phase diagrams of [33, 34]) so to the structural instability. Finally, for a two-dimensional Hamiltonian system one is able to study all its properties in a configuration space rather than in a phase space.

The paper is organized as follows. In Section 2 we present simple Hamiltonian dynamics of the brane universes. In Section 3 we discuss cosmological models in configuration space, stability and the domains of the phase plane for which cosmic acceleration and horizon problem avoidance appear. In Section 4 we present phase portraits for FRW models with dust ($\gamma = 1$) and domain-wall-like matter ($\gamma = 1/3$) [6, 50, 51] which induces accelerated expansion (quintessence) in standard general relativistic cosmology. In section 5 we present simple form of dynamical systems in which constant coefficients play the role of the observational dimensionless density of matter Ω parameters. In Section 6 we discuss our results.

2 Simple dynamics of brane universes

The 5-dimensional Einstein equations for the Randall-Sundrum (RS) model with a brane at $y = 0$ location are [26, 27, 28]

$$\tilde{G}_{\mu\nu}^{(5)} = \kappa_{(5)}^2 [-\Lambda_{(5)} g_{\mu\nu}^{(5)} + \delta(y)(-\lambda h_{\mu\nu}^{(4)} + T_{\mu\nu}^{(4)})] \quad (2)$$

where $\tilde{G}_{\mu\nu}^{(5)}$ is a 5-dimensional Einstein tensor, $g_{\mu\nu}^{(5)}$ is a 5-dimensional metric, $\Lambda_{(5)}$ is a 5-dimensional cosmological constant, and $h_{\mu\nu}^{(4)} = g_{\mu\nu}^{(5)} - n_\mu n_\nu$ is a 4-dimensional induced metric, n^α a unit normal vector to the brane, $T_{\mu\nu}^{(4)}$ – a 4-dimensional energy-momentum tensor, λ is a brane tension, and

$$\kappa_{(5)}^2 = 8\pi^{(5)} G_N = \frac{8\pi}{^{(5)}M_p^3}. \quad (3)$$

where the 5-dimensional Planck mass $^{(5)}M_p$ is always much less than the 4-dimensional $^{(4)}M_p$ as measured on the brane $^{(5)}M_p \ll ^{(4)}M_p = 1.2 \cdot 10^{19} \text{ GeV}$. This fact allows to solve the hierarchy problem in particle physics and possibly reach the electroweak scale $\sim \text{TeV}$ for gravity in accelerators.

The induced 4-dimensional Einstein equations on the brane are [30]

$$G_{\mu\nu}^{(4)} = -\Lambda_{(4)} h_{\mu\nu}^{(4)} + \kappa_{(4)}^2 T_{\mu\nu} + \kappa_{(5)}^2 \Pi_{\mu\nu} - E_{\mu\nu} \quad (4)$$

where

$$\kappa_{(4)}^2 = 8\pi^{(4)} G_N = \frac{8\pi}{^{(4)}M_p^2} = \frac{\lambda}{6} \kappa_{(5)}^4 \quad (5)$$

$$\Lambda_{(4)} = \frac{1}{2} \kappa_{(5)}^2 \left[\Lambda_{(5)} + \frac{1}{6} \kappa_{(5)}^2 \lambda^2 \right] \quad (6)$$

$$\Pi_{\mu\nu} = \frac{1}{12} T T_{\mu\nu} - \frac{1}{4} T_{\mu\alpha} T^\alpha_\nu + \frac{1}{24} g_{\mu\nu} [3T_{\alpha\beta} T^{\alpha\beta} - T^2], \quad (7)$$

$$T_{\mu\nu} = \rho V_\mu V_\nu + p h_{\mu\nu}^{(4)}, \quad (8)$$

where V_μ is the 4-velocity of an observer on the brane, $\Lambda_{(4)}$ is the 4-dimensional cosmological constant on the brane [24, 25] and $E_{\mu\nu}$ is the correction which appears from the Weyl tensor in the bulk which reads as [30, 37, 38]

$$E_{\mu\nu} = -\frac{6\mathcal{U}}{\lambda \kappa_{(4)}^2} \left[V_\mu V_\nu + \frac{1}{3} h_{\mu\nu}^{(4)} + \mathcal{P}_{\mu\nu} + \mathcal{Q}_\mu V_\nu + \mathcal{Q}_\nu V_\mu \right]. \quad (9)$$

Here $\mathcal{U}$ is an effective nonlocal energy density on the brane which arises from the gravitational field in the bulk which is not necessarily positive and reads

$$\mathcal{U} = -\frac{1}{6} \kappa_{(4)}^2 \lambda E_{\mu\nu} V^\mu V^\nu. \quad (10)$$

Since $E_{\mu\nu}$ is traceless, then its effective local pressure is $p = (1/3)\mathcal{U}$. On the other hand, an effective nonlocal anisotropic stress is

$$\mathcal{P}_{\mu\nu} = -\frac{1}{6}\kappa_{(4)}^2\lambda E_{[\mu\nu]}, \quad (11)$$

while an effective energy flux on the brane is

$$\mathcal{Q}_\mu = -\frac{1}{6}\kappa_{(4)}\lambda (E_{\mu\nu}V^\nu + E_{\nu\mu}V^\mu). \quad (12)$$

After admission the perfect fluid with barotropic equation of state $p = (\gamma - 1)\rho$, $\gamma \in [0, 2]$, where p the pressure and ϱ the energy density from (7) one has

$$\Pi_{\mu\nu} = \frac{1}{12}\varrho^2 V_\mu V_\nu + \frac{1}{12}\varrho^2(2\gamma - 1)h_{\mu\nu}^{(4)} \quad (13)$$

and the dynamics of homogeneous models in the RS brane-world scenario can be described by the following set of equations

$$\dot{H} = -H^2 - \frac{(3\gamma - 2)\kappa_{(4)}^2}{6}\rho - \frac{(3\gamma - 1)\kappa_{(4)}^2}{6\lambda}\rho^2 - \frac{2}{3}\sigma^2 + \frac{\Lambda_{(4)}}{3} - \frac{2}{\lambda\kappa_{(4)}^2}\mathcal{U}, \quad (14)$$

$$\dot{\rho} = -3\gamma H\rho, \quad (15)$$

$$\dot{\mathcal{U}} = -4H\mathcal{U}, \quad (16)$$

where $H = d(\ln a)/dt$ is the Hubble function, t is cosmological time, the brane tension is $\lambda > 0$ (this condition allows to recover conventional gravity) and $\kappa_{(4)} = 8\pi G_{(4)}$ is the gravitational coupling constant ($c = 1$). The function $\mathcal{U}$ enters as a contribution from the Weyl tensor in the bulk. The shear scalar $\sigma^2 = \frac{1}{2}\sigma^{ab}\sigma_{ab}$ vanishes for the FRW models, whereas for the anisotropic Bianchi type models

$$\frac{d\sigma^2}{dt} = -6H\sigma^2 \quad \Leftrightarrow \quad \sigma^2 = \sigma_0^2 a^{-6} \quad (17)$$

The first integral of the system (14)–(16) is the Friedmann equation [34] which reads as

$$H^2 = \frac{\kappa_{(4)}^2}{3}\rho + \frac{\kappa_{(4)}^2}{6\lambda}\rho^2 - \frac{k}{a^2} + \frac{\sigma^2}{6} + \frac{\Lambda_{(4)}}{3} + \frac{2\mathcal{U}}{\lambda\kappa_{(4)}^2}, \quad (18)$$

where $k \in \{0, \pm 1\}$ – the curvature index. One can easily see from (18) that in the limit $\lambda \rightarrow \infty$ one recovers general relativity. The main difference from

the standard general relativistic Friedmann equation is the appearance of the ϱ^2 correction. This term comes as a contribution from the brane. However, in the general case within the framework of AdS/CFT correspondence [46] this correction may come from higher-derivative terms in the curvature and higher-dimension operators connecting the on-brane matter directly to the CFT [47].

There are two invariant submanifolds: $\mathcal{U} = 0$ and $\sigma = 0$. The system on the submanifold $\mathcal{U} = 0$ corresponds to dynamics with vanishing electromagnetic part of the Weyl tensor, whereas the system on the submanifold $\sigma = 0$ corresponds to the case without anisotropy (FRW models). The intersection of these submanifolds is also an invariant.

It is well-known that the first integral of the FRW equation can be used to construct a Hamiltonian function. We take advantage of this feature in the considered model.

The integration of (15)–(16) gives

$$\rho = \rho_0 a^{-3\gamma}, \quad \mathcal{U} = \mathcal{U}_0 a^{-4}. \quad (19)$$

Therefore the right-hand side of the Raychaudhuri equation (14) can be expressed in terms of the scale factor $a(t)$ as

$$\ddot{a} = \left[-\frac{(3\gamma - 2)\kappa_{(4)}^2 \rho_0}{6} a^{-3\gamma} - \frac{(3\gamma - 1)\kappa_{(4)}^2 \rho_0^2}{6\lambda} a^{-6\gamma} - \frac{2}{3} \sigma_0 a^{-6} + \frac{\Lambda_{(4)}}{3} - \frac{2\mathcal{U}_0}{\lambda \kappa_{(4)}^2} a^{-4} \right] a. \quad (20)$$

The equation (20) can be rewritten in the form analogous to the Newton equation of motion in the 1-dimensional configuration space $\{a: a \in \mathbb{R}_+\}$

$$\ddot{a} = -\frac{\partial V}{\partial a}, \quad (21)$$

where the potential function

$$V(a) = -\frac{\kappa_{(4)}^2 \rho_0}{6} a^{-3\gamma+2} - \frac{\kappa_{(4)}^2 \rho_0^2}{12\lambda} a^{-6\gamma+2} - \frac{\sigma_0^2}{6} a^{-4} - \frac{\Lambda_{(4)}}{6} a^2 - \frac{\mathcal{U}_0}{\lambda \kappa_{(4)}^2} a^{-2} + V_0, \quad (22)$$

and $V_0 = \text{const.}$ The plot of (22) for $\gamma = 1$ is given in Fig.1. The equation (21) has a simple interpretation: the Randall-Sundrum models *accelerate* whenever the potential $V(a)$ is a decreasing function of a while they *decelerate*

whenever the potential $V(a)$ is an increasing function of a . Another important conclusion from (22) is that for $\gamma > 1$ ($p > 0$) ρ^2 contribution dominates over shear anisotropy near singularity so the singularity in Bianchi models (especially in Bianchi IX) is isotropic and that there is no chaos [38]. In general relativity it is possible only for a meaningless fluid with $\gamma > 2$ ($p > \rho$). The first integral of (21) is

$$V(a) + \frac{\dot{a}^2}{2} = V_0 - \frac{k}{2}. \quad (23)$$

Now we construct the *Hamiltonian function*

$$\mathcal{H} \equiv \frac{\dot{a}^2}{2} + V(a), \quad (24)$$

and then trajectories of the system lie on the energy level $\mathcal{H} \equiv E = \text{const.}$ The advantage of having the Hamiltonian function given by Eq. (24) is that one is then easily able to canonically quantize the system and allow quantum cosmological framework on the brane [39, 40] in full analogy to general relativity [41]. Finally, we obtain the dynamics reduced to the Hamiltonian flow in the 1-dimensional configuration space

$$\mathcal{H} = \frac{\dot{a}^2}{2} + V(a) = 0, \quad (25)$$

with $V(a)$ given by (22) and $V_0 = k/2$.

Now the physical trajectories lie on *the zero-energy level*, $\mathcal{H} = E = 0$, which coincides with the form of the first integral.

The formalism can simply be generalized to the case of a general form of the equation of state. We use this fact and consider the equation of state for the (non-interacting) mixture of dust-like matter ($p = 0$) and an unknown component, labelled X , with negative pressure. It is assumed that dominant X -component is a perfect fluid with the equation of state specified by $p_X = (w_X - 1)\rho_X$ and $0 < w_X < 2/3$, which enables that component to induce accelerated expansion in standard general relativistic cosmology [50, 51]. The positive cosmological constant $\Lambda_{(4)}$ corresponds to $w_X = 0$. The supernovae observations indicate that the dimensionless energy density of X -component $\Omega_{X0} \approx 0.7$ is large compared to dust-like matter ($\rho_{X0} \ll M_{pl}^4$ — natural order of magnitude of the vacuum energy) [6]. The best-fit component is with $w_X \approx 1/3$ (see paper 4 in [6]). One often represents the X -component in terms of a scalar field Φ (e.g., inflaton) with a suitable potential (quintessence) [52, 53]. Conservation of energy gives $\rho_\Phi \propto a^m$, $m = -3w_\Phi$, $p_\Phi = (w_\Phi - 1)\rho_\Phi$, $w_\Phi = \text{const.}$




Figure 1: The potential function $V(a)$ given by (22) for some trajectories depending on the initial conditions in the phase plane for $k = 0, -1, +1$ and $\gamma = 1$ (dust). The horizontal line set by loci formed from maxima of all potential functions separates the regions with *deceleration* (left) and *acceleration* (right) (cf. Eq. (21)). Only the region with $V(a) < 0$ has a physical sense.




Figure 2: The potential function $V(a)$ given by (22) for some trajectories depending on the initial conditions on the phase plane for $k = 0, -1, +1$ and $\gamma = 1/3$ (domain walls) which induces accelerated expansion in standard general relativistic cosmology. Similarly as in Fig. 1 there is the horizontal line of potential maximum points which separates to *deceleration* and *acceleration* regions (cf. Eq. (21)). However it appears for lower values of a .

The most general case of the equation of state is

$$p = p_X + 0 = [\gamma(a) - 1] \rho = \frac{w_X}{\frac{\rho_{m0}}{\rho_{X0}} a^{3(w_X-1)} + 1} \rho, \quad (26)$$

with $w_X = \text{const.}$ and zero stands for the pressure of the dust.

Therefore the total energy $\rho = \rho_m + \rho_X$ changes as

$$\rho = \rho_0 a^{-3} \exp \left(- \int_0^a \frac{3\gamma(a)}{a} da \right). \quad (27)$$

After substitution (26) into (27) we obtain

$$\rho = \rho_{X0} a^{-3w_X} \left[\frac{\rho_{m0}}{\rho_{X0}} a^{3(w_X-1)} + 1 \right] = \rho_{m0} a^{-3} + \rho_{X0} a^{-3w_X}. \quad (28)$$

Then, the potential function is given by the integral

$$V(a) = \int_0^a \left[\frac{(3\gamma(a) - 2)\kappa_{(4)}^2}{6} \rho + \frac{(3\gamma(a) - 1)\kappa_{(4)}^2}{6\lambda} \rho^2 + \frac{2}{3} \sigma^2 - \frac{\Lambda_{(4)}}{3} + \frac{2\mathcal{U}_0}{\lambda\kappa_{(4)}^2} a^{-4} \right] ada, \quad (29)$$

where $\rho(a)$ is given by (28) and $\gamma(a)$ comes from (26).

After substitution the corresponding form of the equation of state (26), the potential $V(a)$ can be obtained by integration of (29)

$$\begin{aligned} V(a) = & -\frac{\kappa_{(4)}^2}{6} \rho_{X0} a^{-3w_X+2} - \frac{\kappa_{(4)}^2}{6} \rho_{m0} a^{-1} \\ & + \frac{\kappa_{(4)}^2}{6\lambda} \left(\rho_{X0} \rho_{m0} a^{-3w_X-1} - \frac{1}{2} \rho_{X0}^2 a^{-6w_X+2} - \frac{1}{2} \rho_{m0}^2 a^{-4} \right) \\ & - \frac{\sigma_0}{6} a^{-4} - \frac{\Lambda_{(4)}}{6} a^2 - \frac{\mathcal{U}_0}{\lambda\kappa_{(4)}^2} a^{-2} + \frac{k}{2}. \end{aligned} \quad (30)$$

It is interesting that in this potential there is a term of the type $\rho_{X0} \rho_{m0} a^{-3w_X-1}$ which is related to the co-existence/interaction of both matter and an unknown form of dark energy in the universe.

Obviously the simplest possibility is the pure case of the unknown matter X such that

$$\rho = \rho_{X0} a^{-3w_X}, \quad p = (w_X - 1)\rho, \quad \gamma(a) = w_X = \text{const},$$

which gives the potential function

$$V(a) = -\frac{\kappa_{(4)}^2}{6} \rho_{X0} a^{-3w_X+2} - \frac{\kappa_{(4)}^2}{12\lambda} \rho_{X0}^2 a^{-6w_X+2} - \frac{\sigma_0^2}{6} a^{-4} - \frac{\Lambda_{(4)}}{6} a^2 - \frac{\mathcal{U}_0}{\lambda\kappa_{(4)}^2} a^{-2} + \frac{k}{2} \quad (31)$$

i.e., w_X plays the role of γ in (22).

As a special example let us consider the case of $p_X = -\frac{2}{3}\rho_X$ ($w_X = \frac{1}{3}$). Then, we have

$$V(a) = -\frac{\kappa_{(4)}^2}{6}\rho_{X0}a - \frac{\kappa_{(4)}^2}{12\lambda}\rho_{X0}^2 - \frac{\sigma_0^2}{6}a^{-4} - \frac{\Lambda_{(4)}}{6}a^2 - \frac{\mathcal{U}_0}{\lambda\kappa_{(4)}^2}a^{-2} + \frac{k}{2}. \quad (32)$$

The diagram of the above function is shown in Fig. 2. Note that the second term in $V(a)$ takes the form of an additive constant like the term $k/2$ and simply scales as curvature term (in general relativity curvature term appears for $w_X = 2/3$ - cosmic strings).

3 Configuration space and stability

In our further analysis we will explore the dynamics given by the canonical equations. Assuming $a = x$ and $y = \dot{a} = \dot{x}$, we have from (25)

$$\dot{x} = \frac{\partial \mathcal{H}}{\partial y} = y \quad (33)$$

$$\dot{y} = -\frac{\partial \mathcal{H}}{\partial x} = -\frac{\partial V}{\partial x}. \quad (34)$$

Then, we can perform a qualitative analysis of the system of autonomous differential equations (33)–(34) in the phase space $(a, \dot{a}) \equiv (x, y)$.

Firstly, we can observe that trajectories are integrable in quadratures. Namely, from the Hamiltonian constraint $\mathcal{H} \equiv E = 0$ we obtain the integral

$$t - t_0 = \int_{a_0}^a \frac{da}{\sqrt{-2V(a)}}. \quad (35)$$

For some specific forms of the potential function (22) we can obviously obtain the exact solutions. However, this is not the task of our paper since we concentrate on qualitative analysis only.

It is possible to make the classification of qualitative evolution paths by analyzing the characteristic curve which represents the boundary equation in the configuration space. For this purpose we consider the equation of zero velocity, $\dot{a} = 0$, which represents the boundary in the configuration space. Because

$$\dot{a}^2 = -2V(a) \quad (36)$$

the motion of the system is limited to the region $\{a: V(a) \leq 0\}$. Consider the boundary of the configuration space given by a condition

$$\partial\mathcal{M} = \{a: V(a) = 0\}. \quad (37)$$

The condition (37) together with (22) is equivalent to the equation of the 4-dimensional cosmological constant on the brane $\Lambda_{(4)}$ expressed as a function of a :

$$\Lambda_{(4)}(a) = \frac{1}{a^2} \left(-\kappa_{(4)}^2 \rho_0 a^{-3\gamma+2} - \frac{\kappa_{(4)}^2 \rho_0^2}{2\lambda} a^{-6\gamma+2} - \sigma_0^2 a^{-4} - \frac{6\mathcal{U}_0}{\lambda \kappa_{(4)}^2} a^{-2} + 3k \right). \quad (38)$$

The plot of $\Lambda_{(4)}(a)$ for different k is shown in Fig. 3 ($\gamma = 1$) and in Fig. 4 ($\gamma = 1/3$). We can see that in the latter case ($\gamma = 1/3$) there is no static Einstein universe. Finally, we consider the evolution path as a level of $\Lambda_{(4)} = \text{const} \neq 0$ and then we classify all evolution modulo their quantitative properties of dynamics (compare [51, 54, 55]). We can conclude that the classes of admissible models in brane scenario are *qualitatively* the same as in general relativity.

The next advantage of representing dynamics in terms of Hamiltonian is the possibility to discuss the stability of critical points which is only based on the geometry of the potential function, namely

- if a diagram of the potential function $V(a)$ has maxima then they correspond to *unstable* critical points; on the other hand, if a diagram of the potential function has minima, they correspond to *stable* attractors.
- if a diagram of the potential function has an inflection point at $a = a_0$, then the corresponding critical point in the phase plane is a *saddle* point.

In general, the stability and the character of a critical point is determined by the Hessian $[\partial^2 \mathcal{H} / \partial x^i \partial y^i]$.

In our case the Hamiltonian function (24) takes the simplest form for natural mechanical systems (i.e., with the kinetic energy quadratic in momenta, and the potential energy dependent on generalized coordinates only). Then, the only possible critical points in a finite domain of phase space are centers and saddles.

The idea of structural stability originated with Andronov and Pontryagin [44]. A dynamical system S is said to be structurally stable if dynamical systems in the space of all dynamical systems are close, in the metric sense, to S or are topologically equivalent to S . Instead of finding and analyzing an individual solution of a model, a space of all possible solutions is investigated. A property is believed to be ‘realistic’ if it can be attributed to large subsets of models within a space of all possible solutions or if it possesses a




Figure 3: The 4-dimensional cosmological constant on the brane, $\Lambda_4(a)$, given by (38), for $k = 0, -1, +1$ and $\gamma = 1$ (dust). Lines $\Lambda_{(4)} = \text{const.}$ gives qualitative classification of possible path evolutions. Let us note that there is a similarity to the standard FRW models. The maximum for $k = +1$ corresponds to a static universe. The de Sitter model starts at the initial singularity. The Eddington model starts from the static universe at $t \rightarrow -\infty$ (separatrices on the phase plane) then reaches the singularity (such that $a(t = 0) = a_0 > 0$) and evolves to infinity. The solutions are represented by levels of constant $\Lambda_{(4)}$ fixed above the $\Lambda_{(4)}(a)$ curve. The domain under the characteristic curve is non-physical.




Figure 4: The 4-dimensional cosmological constant on the brane, $\Lambda_{(4)}(a)$, given by (38), for $k = 0, -1, +1$ and $\gamma = 1/3$ (domain walls). All models with $\Lambda_{(4)}(a) < 0$ and $k = 0, -1$ oscillate. These with $k = +1$ are singularity free. The lines of constant $\Lambda_{(4)}$ above the $\Lambda_{(4)}(a)$ curve represents the qualitative evolution of the model.

certain stability, i.e., if it is shared by a slightly perturbed model. There is a wide opinion among specialists that realistic models should be structurally stable, or even stronger, that everything should possess a kind of structural stability. What does the structural stability mean in physics? The problem is in principle open in more than 2-dimensional case where according to Smale there are large subsets of structurally unstable systems in the space of all dynamical systems [42]. For 2-dimensional dynamical systems, as in the considered case, the Peixoto's theorem says that structurally stable dynamical systems on compact manifolds form open and dense subsets in the space of all dynamical systems on the plane. Therefore, it is reasonable to require the model of a real 2-dimensional problem to be structurally stable.

When we consider the dynamics of brane world models, then there is a simple test of structural stability. Namely, if the right-hand sides of the dynamical systems are in polynomial form, the global phase portraits are structurally stable S^2 ($\mathbb{R}^2$ with a Poincaré sphere) if and only if the number of critical points and limit cycles is finite, each point is hyperbolic and there are no trajectories connecting saddle points. In the considered case the points at infinity are revealed on the projective plane. Two projective maps (z, u) , (v, w) cover a circle at infinity given by $z = 0$ ($z = \infty$) and $v = 0$ ($y = \infty$).

Therefore, one can conclude that the *brane-world models are structurally stable*. It holds because the potential function $V(a)$ is convex up and then there are no non-hyperbolic centres. There are also no trajectories connecting saddle points. All these properties can be deduced from the geometry of the potential function $V(a)$.

Structural stability is sometimes considered as a precondition of the 'real existence'. Having too many drastically different mathematical models - all of them equally well fitting the observational data (up to measurement errors) - seems to be fatal for the empirical method of modern science [45]. Therefore, any 2-dimensional structurally unstable model is not of physical importance.

From the physical point of view it is interesting to answer the question: are the trajectories distributed in the phase space in such a way that critical points are typical or exceptional? How are trajectories with interesting properties distributed? For example, along which trajectories the acceleration condition, $\ddot{a} = -dV/da > 0$, is satisfied? One can easily deduce this from the geometry of the potential function. In the phase space, the *area of acceleration* is determined by $\dot{y} > 0$ or by the condition that

$$\frac{(3\gamma - 2)\kappa_{(4)}^2 \rho_0}{6} a^{-3\gamma} + \frac{(3\gamma - 1)\kappa_{(4)}^2 \rho_0^2}{a}^{-6\gamma} + \frac{2}{3} \sigma_0 a^{-6} - \frac{\Lambda_{(4)}}{3} a + \frac{2\mathcal{U}_0}{\lambda \kappa_{(4)}^2} a^{-4} < 0. \quad (39)$$

From (39) we can see that in order to obtain acceleration, the *positive* 4-dimensional cosmological constant $\Lambda_{(4)}$ is necessary.

It can easily be demonstrated that if $\dot{a}(t) \rightarrow \text{const}$ as $a \rightarrow 0$, then the corresponding world model has *no particle horizon*. Indeed, if there exists a constant C such that for sufficiently large ϵ , $da/dt < C$, then

$$\int_0^{a_0} \frac{da}{a} < C \int_0^{t_0} \frac{dt}{a} = C(\eta_0 - \eta_{\text{sing}}). \quad (40)$$

The integral on the left hand-side of this formula diverges which means that the time η goes to minus infinity, and that there are no causally disconnected regions. Putting this in terms of variables $\{x, y\}$ one needs $x \rightarrow 0$, $y \rightarrow \text{const}$ for the horizon problem to be solved. Now, from the Hamiltonian constraint we obtain that as $x \rightarrow 0$, then $V(x) \rightarrow \text{const}$. Therefore, if the *horizon problem is solved*, then the curvature effects cannot be neglected in the vicinity of an initial singularity.

4 Phase plane analysis of the models with $\gamma = 1$ and $\gamma = 1/3$

In this section we discuss the dynamics of the brane universes in a more detailed way. Firstly, it is easy to verify that the general relativistic limit can be recovered if one takes

$$\frac{1}{\lambda} \rightarrow 0$$

in Eq. (18). The quadratic contribution of the brane ρ^2 matters only when [31]

$$\rho > \lambda > (100 \text{ GeV})^4.$$

In order to simplify Eq. (18) one can assume that $\kappa_{(4)}^2 = 1$ or ${}^{(4)}M_p^2 = 1/8\pi = (1.2 \times 10^{19} \text{ GeV})^2$ which gives

$$\lambda = \frac{6}{\kappa_5^4}, \quad \Lambda_{(4)} = \frac{1}{2}\kappa_5^2\Lambda_{(5)} + \frac{1}{2}\lambda = \frac{1}{2}\sqrt{\frac{6}{\lambda}}\Lambda_{(5)} + \frac{1}{2}\lambda$$

and finally Eq. (18) reads as

$$H^2 = \frac{\rho}{3} \left(1 + \frac{\rho}{2\lambda}\right) + \frac{1}{6} \left(\sqrt{\frac{6}{\lambda}}\Lambda_{(5)} + \frac{1}{2}\lambda\right) + \frac{\sigma_0^2}{6}a^6 - ka^{-2} + \frac{2\mathcal{U}_0}{\lambda}a^{-4}. \quad (41)$$

Therefore, the dynamics depends on four independent constants. Without losing a generality, we put $\Lambda_{(5)} = \mathcal{U}_0 = \rho_0 = 1$. Finally, the dynamics depends on one parameter λ (brane tension) only.

As a proof of the effectiveness of the presented method we consider only two limit subcases: pure dust matter and pure matter X .

Then, the dynamical system (33)-(34) has the form

$$\dot{x} = y \quad (42)$$

$$\dot{y} = -\frac{3\gamma-2}{6}x^{-3\gamma+1} - \frac{3\gamma-1}{6\lambda}x^{-6\gamma+1} - \frac{2}{3}\sigma_0x^{-5} + \frac{\Lambda_{(4)}}{3}x - \frac{2}{\lambda}x^{-3} \quad (43)$$

where $\Lambda_{(4)} = \frac{1}{2}\sqrt{\frac{6}{\lambda}} + \frac{1}{2}\lambda$. We can see that the vector field in (43) is not smooth. There are two important cases: $\sigma_0 = 1$ and $\sigma_0 = 0$. The former describe the Bianchi models (Bianchi I for $k = 0$ and Bianchi V for $k = -1$). The latter describes the FRW models ($k = 0, \pm 1$). The qualitative dynamics of both is similar. Due to the existence of the energy integral (25), the phase space is separated into the two domains by the flat model trajectory (Fig. 5).

Let us consider first the case of the FRW dynamics with dust $\gamma = 1$. In this case the effects of brane and shear in the Bianchi I models are equivalent. An interesting result which shows that there exists a maximum of scalar shear in the Bianchi I model was obtained by Toporensky [49].

The system (42)-(43) can be regularized at the origin, after introducing the projective coordinates

$$z = \frac{1}{x}, \quad u = \frac{y}{x}. \quad (44)$$

Then, we obtain

$$\dot{z} = -uz \quad (45)$$

$$\dot{u} = -\frac{1}{6}z^3 - \frac{2}{3\lambda}z^6 + \frac{\Lambda_{(4)}}{3} - \frac{2}{\lambda}z^4 - u^2. \quad (46)$$

The system has only three critical points in this map

$$\begin{aligned} u_0 &= \pm\sqrt{\frac{\Lambda_4}{3}}, & z_0 &= 0, \\ u_0 &= 0, & z &= z_0: 2z^6 + 12z^4 + \lambda z^3 - 2\Lambda_{(4)}\lambda = 0. \end{aligned}$$

In order to find that there is one critical point with $z_0 > 0$, it is sufficient to consider the diagram of the function

$$f(z) = 2z^6 + 12z^4 + \lambda z^3 - 2\Lambda_{(4)}\lambda.$$




Figure 5: The division of phase space for system (42)-(43) on different domains with respect to the curvature index. The flat model trajectory $k = 0$ separates the regions of the models with negative and positive curvature.




Figure 6: The phase portrait of the system (42)–(43) with $\lambda = 10^8$ and $\gamma = 1$. The Einstein-de Sitter and Eddington models are represented by separatrices in the phase plane. In the neighbourhood of separatrices one can see the Lemaître-Eddington (L-E) evolution with characteristic quasi-static regime. The closer to a critical point the trajectory is, the longer the time of a quasi-static stage evolution. All phase curves lie on algebraic curves given by the first integral (Hamiltonian constraint). The acceleration region is situated to the right of the saddle point, therefore for the L-E universes the acceleration begins in the middle of quasi-static stage.




Figure 7: The phase portrait of system (42)–(43) with $\lambda = 10^8$ and $\gamma = 1/3$. The negative curvature term and the brane term in potential (22) are of the same type. There are two types of trajectories which expand to the maximal scale factor a and then recollapse or vice versa. In the first type of evolution the closed models start with the singularity $a(0) = 0$ and $\dot{a} = \infty$, reach the maximum radius and recollapse while the other class of models recollapse from $a(0) = \infty$ and $\dot{a} = -\infty$ to the minimum value of a , and then expand to infinity. As in the $\gamma = 1$ case, the acceleration region is on the right to the saddle point.

If $\Lambda_{(4)}\lambda > 0$ then $f(0) < 0$ and because $f(z)$ is strictly increasing as $z > 0$ then there exists only one critical point such that $z_0 > 0$, such that $f(z_0) = 0$. Let us note that for $\Lambda_{(4)} < 0$ there is no such a point. The phase portrait of this system is presented in Fig. 8.

For the completeness of the analysis of the dynamical system, it is necessary to consider another projective map (v, w)

$$v = \frac{1}{y}, \quad w = \frac{x}{y}. \quad (47)$$

After regularization and after introducing a new time variable $\eta: d\eta = dt/w^5$ we obtain from (42)-(43) and (47)

$$\dot{v} = v \left(\frac{1}{6}v^3w^3 + \frac{1}{3\lambda}v^6 - \frac{\Lambda_{(4)}}{3}w^6 + \frac{2}{\lambda}v^4w^2 \right) \quad (48)$$

$$\dot{w} = w \left(w^4 + \frac{1}{6}v^3w^3 + \frac{1}{3\lambda}v^6 - \frac{\Lambda_{(4)}}{3}w^6 + \frac{2}{\lambda}v^4w^2 \right). \quad (49)$$

In this map we can find only one critical point $v = 0$ and $w = 0$. As it was mentioned before, it is interesting to observe how much time the trajectories spend in the area of accelerated expansion defined as

$$\{(x, y): x > 0 \wedge \dot{y} > 0\} \Leftrightarrow \left\{ (x, y): \frac{3\gamma - 2}{6}x^{-3\gamma} + \frac{3\gamma - 1}{6\lambda}x^{-6\gamma} + \frac{2}{3}\sigma_0x^{-6} - \frac{\Lambda_{(4)}}{3} + \frac{2}{\lambda}x^{-4} < 0 \right\}.$$

For the special case of dust we obtain in (z, u) -coordinates

$$D_{\text{accel}} = \left\{ (z, u): \frac{1}{6}z^3 + \frac{1}{3\lambda}z^6 + \frac{2}{3}\sigma_0z^6 - \frac{\Lambda_{(4)}}{3} + \frac{2}{\lambda}z^4 > 0 \wedge z > 0 \right\}.$$

The dynamical system for pure accelerating matter has the form analogous to (42)-(43) but now w_X coincides to γ . As an illustration we consider the case of $p_X = -\frac{2}{3}\rho_X$ (domain-wall-like matter), i.e. $w_X = \frac{1}{3}$. Then, we have

$$\dot{x} = y \quad (50)$$

$$\dot{y} = \frac{1}{6} - \frac{2}{\lambda}x^{-3} + \frac{\Lambda_{(4)}}{3}x. \quad (51)$$

In this case, the term which comes from the brane and which is proportional to ρ^2 vanishes (see Eq. 18), and only the term which comes from a non-vanishing Weyl tensor is present in (50)-(51).




Figure 8: The phase portrait of system (45)–(46) with $\lambda = 10^8$ and $\gamma = 1$. All closed generic models are of two types. The first starts from the anti-de Sitter and finishes at de Sitter stage. The second starts from an initial singularity, expands to a maximum size, and then recollapses to a second singularity. Non-generic (exceptional) cases corresponds to the separatrices going in or out of the saddle point, which represent the static universe. For all open and expanding universes the de Sitter model is a global attractor, and for open and contracting universes the anti-de Sitter is a global repeller. The singularity is reached at $z = \infty$ and the half-plane $z < 0$ has no physical sense. All points at the infinity which are represented by $\{z = 0\}$ axis are hyperbolic, therefore the system is *structurally stable*. The same result is valid in the (v, w) plane. The brane effects produce ρ^2/λ term which is formally equivalent to the effects of the stiff matter equation $p = \rho$ (or a massless scalar field).




Figure 9: The phase portrait of system (52)–(53) with $\lambda = 10^8$ and $\gamma = 1/3$. The acceleration region is situated on the left from the saddle point. Therefore the recollapsing and then expanding models lie permanently in the acceleration domain. The Eddington model is also in this region. The Lemaître-Eddington type models start accelerating in the middle of quasi-static phase.

After introducing the projective coordinates

$$z = \frac{1}{x}, \quad u = \frac{y}{x}$$

we obtain

$$\dot{z} = -uz \tag{52}$$

$$\dot{u} = \frac{1}{6}z - \frac{2}{\lambda}z^4 + \frac{\Lambda_{(4)}}{3} - u^2. \tag{53}$$

The critical points are

$$\begin{aligned} u_0 &= \pm \sqrt{\frac{\Lambda_{(4)}}{3}}, & z_0 &= 0, \\ u_0 &= 0, & z_0 &= z: f(z) = \frac{1}{6}z - \frac{2}{\lambda}z^4 + \frac{\Lambda_{(4)}}{3} = 0. \end{aligned}$$

It can be shown that $f(z_0) = 0$ always exists because $f(0) > 0$ and $f(\infty) = -\infty$.

In turn, in the map

$$v = \frac{1}{y}, \quad w = \frac{x}{y}$$

we have

$$\frac{dv}{d\eta} = w^3 \frac{dv}{dt} = -v \left(\frac{1}{6}vw^3 + \frac{\Lambda_{(4)}}{3}w^4 - \frac{2}{\lambda}v^4 \right) \tag{54}$$

$$\frac{dw}{d\eta} = w^3 \frac{dw}{dt} = w^3 - w \left(\frac{1}{6}vw^3 + \frac{\Lambda_{(4)}}{3}w^4 - \frac{2}{\lambda}v^4 \right). \tag{55}$$

From (50)–(51) one can easily determine the domain of acceleration which corresponds to the domain of x in which $V(x)$ is a decreasing function of its argument

$$-\frac{dV}{dx} > 0 \quad \Leftrightarrow \quad x > x_{\max}$$

where $x_{\max} : \frac{\partial V}{\partial x}|_{x=x_{\max}} = 0$ or $\dot{y}(x_{\max}) = 0$. It can be checked that $\dot{y}$ is a strictly increasing function of a ($\ddot{y} > 0$) and $\dot{y}(0) = -\infty$, so there is a single point $x_{\max}$.

From (51) we can also observe that as x goes to zero then y goes to infinity, which means that the model has a particle horizon ($\Lambda_{(4)}$ is negligible and then $y \propto x^{-1}$). It is due to the presence of $u_0 \neq 0$.

From (42)–(43) and the integral of energy we can see that we have the horizon in a generic case if $\rho_0 \neq 0 \wedge u_0 \neq 0 \wedge \sigma_0 \neq 0$. In the special case of $u_0 = \sigma_0 = 0$ we have a model without a horizon (the FRW models with vanishing electromagnetic part of Weyl tensor).

From (50)–(51), we obtain that as $x \rightarrow 0$, then $\dot{y} = \frac{1}{6}$, i.e. $y \propto t$ and $x(t) \propto t^2$.

In general, from the first integral (25) one observes that the generic evolution of cosmological models without a horizon is when $V(a) \rightarrow \text{const.}$ (zero is also possible) as $a \rightarrow 0$.

Effects of $\Lambda_{(4)}$ are always negligible near a singularity and if $u_0 \neq 0$ we have always asymptotically $V(0) = \infty$. For $u_0 = \sigma_0 = 0$ and $\gamma \leq \frac{1}{3}$ (domain walls on the brane), then $V(a)$ goes to a constant and there is no horizon in the model. Then, the generic behaviour near the singularity is $x(t) \propto t$, i.e., it is Milne's evolution (in general relativity this appears for $\gamma = 2/3$ (cosmic strings on the brane [51])).

5 Simple dynamics in terms of density parameters Ω and the exact solutions

Independent observations of supernovae type Ia, made by the Supernovae Cosmology Project and the High z Survey Team [6], indicate that our Universe is currently accelerating. There is a fundamental problem for theoretical physics to explain the origin of this acceleration. If we introduce the cosmological constant $\Lambda_{(4)}$ and assume that the Universe is flat, then the best-fit model is for the cosmological constant density parameter $\Omega_{\Lambda_{(4)},0} = 0.72$ and for the dust density parameter equal to $\Omega_{m,0} = 0.28$ (index "0" refers to the present moment of time).

Our formalism gives natural base to express dynamical equations in terms of dimensionless observational density parameters Ω and to compare the results with supernovae data. However, before we study these quantities in detail following the discussion of Refs. [51, 55, 56, 58] we introduce the notation in which it is easy to tell which models are exactly integrable. The Friedmann Eq. (18) with the help of the conservation equations (19) and (20), can be rewritten in the form

$$\frac{1}{a^2} \left(\frac{da}{dt} \right)^2 = \frac{C_{GR}}{a^{3\gamma}} + \frac{C_\lambda}{a^{6\gamma}} - \frac{k}{a^2} + \frac{\Lambda_4}{3} + \frac{C_U}{a^4}, \quad (56)$$

where we have defined the appropriate constants ($\kappa_{(4)}^2 = 8\pi G$)

$$C_{GR} = \frac{\kappa_{(4)}^2}{3} a^{3\gamma} \varrho, \quad (57)$$

$$C_\lambda = \frac{\kappa_{(4)}^2}{6\lambda} a^{6\gamma} \varrho^2, \quad (58)$$

$$C_{\mathcal{U}} = \frac{2}{\kappa_{(4)}^2 \lambda} a^4 \mathcal{U}, \quad (59)$$

and C_{GR} is of a general relativistic nature. It is easy to notice that the following cases can be exactly integrable in terms of elliptic functions [55, 51]: $\gamma = 0$ (cosmological constant), $\gamma = 1/3$ (domain-wall-like matter) and $\gamma = 2/3$ (cosmic strings). The first case is the easiest since in this case the first two terms on the right-hand-side of (56) play the role of cosmological constants similar to $\Lambda_{(4)}$. The next two cases involve terms which were already integrated in the context of general relativity. For $\gamma = 1/3$ (domain-wall-like matter on the brane) the general relativistic term with C_{GR} in (56) scales as domain-walls in general relativity while the term with C_λ scales as cosmic string (curvature) in general relativity. For $\gamma = 2/3$ the term with C_λ scales as radiation in general relativity. Then, the problem of writing exact solutions reduces to the repetition of the discussion of Refs. [55, 51]. We will not be doing this here. For other values of $\gamma = 4/3; 1; 2$ the terms of the type $1/a^8$ and $1/a^{12}$ appear and the integration involves hyperelliptic integrals.

Coming back to observational quantities we now define the dimensionless observational density parameters [56, 57, 58]

$$\Omega_{GR} = \frac{\kappa_{(4)}^2}{3H^2} \varrho, \quad (60)$$

$$\Omega_\lambda = \frac{\kappa_{(4)}^2}{6H^2 \lambda} \varrho^2, \quad (61)$$

$$\Omega_{\mathcal{U}} = \frac{2}{\kappa_{(4)}^2 H^2 \lambda} \mathcal{U}, \quad (62)$$

$$\Omega_k = -\frac{k}{H^2 a^2}, \quad (63)$$

$$\Omega_{\Lambda_{(4)}} = \frac{\Lambda_{(4)}}{3H^2}, \quad (64)$$

where the Hubble parameter and the deceleration parameter read as

$$H = \frac{\dot{a}}{a}, \quad (65)$$

$$q = -\frac{\ddot{a}a}{\dot{a}^2}, \quad (66)$$

so that the Friedmann equation (18) can be written down in the form

$$\Omega_{GR} + \Omega_{\lambda} + \Omega_k + \Omega_{\Lambda_{(4)}} + \Omega_{\mathcal{U}} = 1. \quad (67)$$

Using (60)-(66) the equation (20) can now be rewritten as (compare Eq.(10) of [56])

$$\frac{\Lambda_{(4)}}{3H^2} = \frac{3\gamma - 2}{2}\Omega_{GR} + \frac{3\gamma - 1}{2}\Omega_{\lambda} + \Omega_{\mathcal{U}} - q. \quad (68)$$

On the other hand, it is useful to express the curvature of spatial sections by observational quantities using (67) and (68)

$$\frac{k}{H^2 a^2} = \frac{3\gamma}{2}\Omega_{GR} + \frac{3\gamma + 1}{2}\Omega_{\lambda} + 2\Omega_{\mathcal{U}} - q - 1. \quad (69)$$

These relations (68) and (69) together with the exact solutions (which here are certainly available for $\gamma = 0; 1/3; 2/3$) (see Refs. [58, 59]) are useful in writing down an explicit redshift-magnitude formula (generalized Hubble law) for the brane models to study their compatibility with astronomical data from supernovae. Such a comparison will be presented elsewhere [60].

In this paper we only study the formalism which allows to formulate our phase space quantities in terms of the observational parameters Ω . Let us consider a brane universe filled with an unknown type of matter with barotropic equation of state $p_i = (w_i - 1)\rho_i$. Then, it is useful to rewrite the dynamical equations to a new form using dimensionless quantities

$$x \equiv \frac{a}{a_0}, \quad T \equiv |H_0|t. \quad (70)$$

The basic dynamical equations are then rewritten as

$$\frac{\dot{x}^2}{2} = \frac{1}{2}\Omega_{k,0} + \frac{1}{2}\sum_i \Omega_{i,0}x^{2-3w_i} = -V(x) \quad (71)$$

$$\ddot{x} = \frac{1}{2}\sum_i \Omega_{i,0}(2 - 3w_i)x^{1-3w_i}, \quad (72)$$

where $\Omega_i = (\Omega_{GR}, \Omega_{\lambda}, \Omega_{\mathcal{U}}, \Omega_{\Lambda_4})$.

The above equations can be represented as a two-dimensional dynamical system

$$\begin{aligned} \dot{x} &= y \\ \dot{y} &= \frac{1}{2}\sum_i \Omega_{i,0}(2 - 3w_i)x^{1-3w_i}. \end{aligned}$$

As we have mentioned already the dynamics can be always reduced to the form of the Friedmann models of general relativity with a certain kind of a non-interacting multifluid.

Of course, the dynamical system (71)-(72) has the Hamiltonian

$$\mathcal{H} = \frac{p_x^2}{2} + V(x), \quad (73)$$

where $V(x) = -\frac{1}{2}\Omega_{k,0} - \frac{1}{2}\sum_i \Omega_{i,0}x^{2-3w_i}$, which should be considered on the zero-energy level.

As an example of application of these equations to study the problem of cosmic acceleration, consider the case of $\Omega_{\mathcal{U},0} = 0$ and $\Omega_{\Lambda_{(4)},0} \neq 0$. It emerges that at present our Universe accelerates provided that

$$\Omega_{X,0}(2 - 3w_X)x^{1-3w_X} + \Omega_{\lambda,0}(2 - 6w_X)x^{1-6w_X} - 2\Omega_{\Lambda_{(4)},0}x > 0, \quad (74)$$

where w_X refers to any type of matter X with the equation of state $p_X = (w_X - 1)\varrho_X$ (for $\Omega_{\Lambda_{(4)}} = 0$ it is possible if $w_X < 1/3$).

It is convenient to introduce a new variable $z = x^{-3w_X} > 0$, then Eq. (74) reduces to the quadratic inequality

$$\Omega_{\lambda,0}(2 - 6w_X)z^2 + \Omega_{X,0}(2 - 3w_X)z - 2\Omega_{\Lambda_{(4)},0} > 0. \quad (75)$$

In the special case of the component X in the form of the wall-like-matter $w_X = 1/3$, the Universe accelerates if

$$\Omega_{X,0}z > -2\Omega_{\Lambda_{(4)},0}, \quad (76)$$

i.e., if $\Lambda_{(4)} > 0$, $w_X > 0$, the universe always accelerates.

In general, the domain of acceleration depends on the solution of the inequality (75). We assume that we are in the region of acceleration $\Omega_{\Lambda_{(4)},0} > 0$, $w_X > 0$ and consider two cases.

Case 1: $w_X > 1/3$.

There are always two solutions for z with opposite signs. One solution is for $0 < z < z_+$ (or $x_+ < x < \infty$), where x_+ is given by

$$x_+ = \left[\frac{(3w_X - 2)\Omega_{X,0} + \sqrt{[(2 - 3w_X)\Omega_{X,0}]^2 - 16(1 - 3w_X)\Omega_{\Lambda_{(4)},0}\Omega_{\lambda,0}}}{(3w_X - 1)\Omega_{\lambda,0}} \right]^{-1/3w_X}. \quad (77)$$

In this case we have a minimum value of acceleration at

$$x_{\min} = \left[\frac{(2 - 3w_X)\Omega_{X,0}}{2(6w_X - 2)\Omega_{\lambda,0}} \right]^{-1/3w_X}. \quad (78)$$

Case 2: $w_X < 1/3$.

The universe accelerates for every x .

Let us note that in Case 1 the minimum value of x can always be expressed in terms of redshift $z = x^{-1} - 1$ and from the inequality $x_+ \leq 1$ we obtain an additional restriction on parameters $\Omega_{i,0}$.

6 Conclusion

In the paper we studied the dynamics of Randall-Sundrum brane universes with isotropic Friedmann geometry. Our approach is the simplest in that we formulate the problem in a 2-dimensional phase space and not in a higher-dimensional phase space as it has been done recently. First of all, this is the smallest possible dimension to study the isotropic cosmological systems of equations. Then, it allows to *avoid* the degeneracy of critical points and trajectories which appear in a higher dimensional phase space and so to *avoid* structural instability of these models. Also, since the system is Hamiltonian one is able to study its properties in a one-dimensional configuration space rather than in phase space. In such a space the motion of the universe point is reduced to the motion of a particle with the potential energy $V(a)$, where a is the scale factor (see Section 2). On the other hand, Hamiltonian formulation can easily be applied to quantum cosmology on the brane.

In Fig. 1 and 2 we can see that there is no qualitative difference in the shapes of the potential function $V(a)$ as γ varies for cases under studies. Only the values of a for which the potential is negative have the physical meaning. The maximum of the potential function corresponds to an unstable saddle point on a phase diagram (a critical point solution of the zero energy level). The de Sitter solution is a point at infinity. Near the initial singularity we have a solution $a(t) \propto t^{1/3\gamma}$ which differs from a general relativistic one $a(t) \propto t^{2/3\gamma}$.

From the theory of qualitative differential equations in Sections 3 and 4 we obtained the visualization of the system evolution in the phase plane $(x, \dot{x})$ and analyzed the asymptotic states and concluded that the brane models are *structurally stable*. We studied the solution of the (past) horizon problem and the initial conditions for acceleration of our Universe. We have the neat interpretation of a *domain of acceleration* as a domain in configuration space where the potential function decreases. Therefore from the observation of the potential function $V(a)$, we can see the acceleration domain $a > a_{\max}$, with $V'(a_{\max}) = 0$, which is independent of the curvature index k . On the other hand, the (past) horizon problem is solved when $V(a) \rightarrow \text{const}$ (may be equal to zero) as $a \rightarrow 0$. If we find trajectories for which $y = \dot{a}$ goes to

infinity as $x = a \rightarrow 0$, then the horizon is present in such a model. This is the case of dust brane model as well as a brane world filled with domain-wall-like matter with $\gamma = 1/3$.

In general our model does not solve the horizon problem due to the existence of matter dominated phase of the early evolution where the term ρ^2 dominates and so $H \propto \rho \propto a^{-3\gamma}$ ($a(t) \propto t^{-1/3\gamma}$). Only in the special case of heavy domain-wall-like matter ($\gamma \leq 1/3$) the horizon problem is solved, because the evolution near the singularity dominated by matter (brane tension) is $\dot{a} \propto a^{1-3\gamma}$. However, the existence of such a singularity requires $\mathcal{U} = 0$ (Weyl tensor contribution from the bulk vanishes). Therefore, the case $\gamma \leq 1/3$ is strictly distinguished because as $x \rightarrow 0$ then $y \rightarrow \text{const}$. For $\gamma = 1/3$ the term ρ^2/λ vanishes and brane effects are negligible.

In the phase portraits we can observe similarities as well as differences in both considered cases. In the dust-like matter case there is a quasi-static stage first discussed by Lemaître (called ‘loitering stage’ in [48]). Such quasi-static stages, present in the case of wall-like-matter, corresponds to configurations in the vicinity of a critical point. The acceleration does not depend on the value of Ω_k , but in the phase space there are different sets of initial conditions which provide acceleration.

We conclude that the classes of admissible models in brane scenario are *qualitatively* the same as in general relativity, i.e., there exists a homeomorphism in the phase space which transforms trajectories of brane models into general relativistic ones without a change of orientation. However, *quantitatively* the models differ from general relativistic ones - for instance, they asymptote the singularity in a different way.

Finally, we reformulated the phase space quantities in terms of the observational parameters Ω and presented the formalism in which one is able to perform the observational tests (redshift-magnitude relation) of brane models which we will address in a separate paper.

7 Acknowledgements

M.S. and A.K. were partially supported by the Polish Research Committee (KBN) Grant No 2PO3B 107 22.

References

- [1] G. Veneziano, Phys. Lett. **B265** (1991), 287.
- [2] M. Gasperini and G. Veneziano, Astropart. Phys. **1** (1993), 317.

- [3] J.E. Lidsey, D.W. Wands, and E.J. Copeland, Phys. Rep. **337** (2000), 343.
- [4] A.H. Guth, Phys. Rev. D**23** (1981), 347.
- [5] A.D. Linde, Phys. Lett. **108** (1982), 389.
- [6] S. Perlmutter *et al.*, Nature **391**, (1998) 51; S. Perlmutter *et al.*, Ap. J. **517**, (1999) 565; P. M. Garnavich *et al.* Ap. J. Lett. **493** (1998) L53; A. G. Riess *et al.* Astron. J. **116** (1998) 1009.
- [7] R.R. Caldwell, R. Dave and P.J. Steinhardt, Phys. Rev. Lett. **80** (1998), 1582; I. Zlatev, L. Wang and P.J. Steinhardt, Phys. Rev. Lett. **82** (1999), 896; P.J. Steinhardt, L. Wang and I. Zlatev, Phys. Rev. D**59** (1999), 123504.
- [8] P. Hořava and E. Witten, Nucl. Phys. B**460** (1996), 506; *ibid* B**475**, 94.
- [9] L. Randall and R. Sundrum, Phys. Rev. Lett., **83** (1999), 3370.
- [10] L. Randall and R. Sundrum, Phys. Rev. Lett., **83** (1999), 4690.
- [11] R.Y. Donagi, J. Khoury, B. Ovrut, P.J. Steinhardt and N.J. Turok, JHEP **0111** (2001), 041.
- [12] J. Khoury, B. Ovrut, N. Seiberg, P.J. Steinhardt and N.J. Turok, Phys. Rev. D**65** (2002), 086007.
- [13] E. Witten, *The Cosmological Constant from the Viewpoint of String Theory*, e-print: hep-ph/0002297; *Quantum Gravity in De Sitter Space*, e-print: hep-th/0106109.
- [14] S. Hellerman, N. Kaloper and L. Susskind JHEP **0106** (2001), 003; W. Fishler, A. Kashani-Poor, R. McNees, S. Paban JHEP **0107** (2001), 003; J.W. Moffat *Acceleration of the Universe, String Theory and a Varying Speed of Light*, e-print: hep-th/0105017.
- [15] M. Visser, Phys. Lett. B**159** (1985), 22.
- [16] C. Barcelo and M. Visser, Phys. Lett. B**482** (2000) 183.
- [17] A. Lukas, B.A. Ovrut and D. Waldram, Phys. Rev. D**60** (1999), 086001.
- [18] H.S. Reall, Phys. Rev. D**59** (1999), 103506.
- [19] J.E. Lidsey, Class. Quantum Grav., **17** (2000), L39.

- [20] M.P. Dąbrowski, Phys. Lett. B**474** (2000), 52.
- [21] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, Phys. Lett. B**516** (1998) 70.
- [22] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, G. Dvali, Phys. Lett. B**436** (1998) 257.
- [23] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, Phys. Rev. D**59** (1999) 086004.
- [24] P. Binétruy, C. Deffayet and D. Langlois, Nucl. Phys. B**565** (2000), 269.
- [25] P. Binétruy, C. Deffayet and D. Langlois, Phys. Lett. B**477** (2000), 285.
- [26] T. Shiromizu, K. Maeda, and M. Sasaki, Phys. Rev. D**62**, 024012 (2000).
- [27] M. Sasaki, T. Shiromizu and K. Maeda, Phys. Rev. D**62** (2000), 024008.
- [28] S. Mukhoyama, T. Shiromizu and K. Maeda, Phys. Rev. D**62**, 024028.
- [29] H. Ishihara, Phys. Rev. Lett. **86** (2001), 381.
- [30] R. Maartens, *Geometry and dynamics of the brane-world*, e-print:gr-qc/0101059.
- [31] L.A. Gergely and R. Maartens, Class. Quantum Grav. **19** (2002) 213.
- [32] A. Frolov, Phys. Lett. B**514** (2001) 213.
- [33] A. Campos and C.F. Sopuerta, Phys. Rev. D**63** (2001), 104012.
- [34] A. Campos and C. F. Sopuerta, Phys. Rev. D**64** (2001), 104011.
- [35] C.M. Chen, T. Harko and M.K. Mak, Phys. Rev. D**64** (2001), 044013.
- [36] C.M. Chen, T. Harko and M.K. Mak, Phys. Rev. D**64** (2001), 124017.
- [37] A. A. Coley, *Dynamics of brane-world cosmological models*, e-print: hep-th/0110049.
- [38] A. A. Coley, Class. Quant. Grav. **19** (2002) L45.
- [39] L. Anchordoqui, C. Nuñez and K. Olsen, JHEP 0010:050 (2000).
- [40] L. Anchordoqui *et al.*, Phys. Rev. D**64** (2001) 084027.
- [41] M.P. Dąbrowski and A.L. Larsen, Phys. Rev. D**55** (1995), 3424.

- [42] S. Smale, *Mathematics of Time* (Springer Verlag, New York, 1980).
- [43] O. I. Bogoyavlensky, *Methods in the Qualitative Theory of Dynamical Systems in Astrophysics and Gas Dynamics* (Springer-Verlag, Berlin, 1985).
- [44] A. A. Andronov and L. S. Pontryagin, Systèmes grossiers. Dokl. Akad. Nauk SSSR **14** (1937), 247–250.
- [45] R. Thom, *Stabilité structurelle et morphogénese* (Inter-Editions, Paris, 1997).
- [46] J. Maldacena, Adv. Theor. Math. Phys. **2** (1998), 231.
- [47] A. Hebecker, J. March-Russell, Nucl. Phys. B**608** (2001), 375.
- [48] V. Sahni, H. Feldman and A. Stebbins, Ap. J. **385** (1992), 1.
- [49] A. V. Toporensky, Class. Quant. Grav. **18** (2001), 2311.
- [50] J. Stelmach, M. P. Dąbrowski and R. Byrka, *Nuclear Physics* **B406**, (1993), 471.
- [51] M.P. Dąbrowski, Ann. Phys (N. Y.) **248** (1996) 199.
- [52] K.I. Maeda, Phys. Rev. D**64** (2001), 123525.
- [53] P.F. González-Díaz, Phys. Lett. B**481** (2000), 353.
- [54] H. P. Robertson, Rev. Mod. Phys. **5** (1933) 62.
- [55] M.P. Dąbrowski and J. Stelmach, Ann. Phys (N. Y.) **166** (1986) 422.
- [56] M.P. Dąbrowski and J. Stelmach, Astron. Journ. **92** (1986), 1272.
- [57] M.P. Dąbrowski and J. Stelmach, Astron. Journ. **93** (1987), 1373.
- [58] M.P. Dąbrowski and J. Stelmach, Astron. Journ. **92** (1986), 1272.
- [59] M. P. Dąbrowski and J. Stelmach, in *Large Scale Structure in the Universe* IAU Symposium No 130 eds. Jean Adouze, Marie-Christine Palletan and Alex Szalay, (Kluwer Academic Publishers, 1988), 566.
- [60] M. Szydłowski, M. P. Dąbrowski and A. Krawiec, in preparation.